Research letter

A case of rilpivirine drug-induced liver injury

Liver toxicity is a frequent adverse event following antiretroviral therapy for HIV, ranging from asymptomatic increases in liver enzymes to fulminant liver failure.^{1,2} Rilpivirine is a second-generation non-nucleoside reverse transcriptase inhibitor with low rates of liver enzyme elevations.³ We report a case of acute hepatitis secondary to rilpivirine with histological documentation.

A 27-year-old Mediterranean man was diagnosed with HIV, with a baseline CD4 count of $0.359 \times 10^9/L$ and an HIV-1 viral load of 21 525 copies/mL. His last negative HIV test was 2 years prior. His medical history included depression, G6PD deficiency and a beta thalassaemia carrier status. He reported minimal alcohol intake. His baseline liver markers were normal.

Abacavir/lamivudine 600/300 mg once daily and raltegravir 400 mg two times per day were started on week 4 following diagnosis. Raltegravir was switched to rilpivirine 25 mg once daily on week 11 due to symptoms of fatigue.

His liver markers were abnormal on week 25, 14 weeks following switch, with alanine transaminase (ALT) of 231 IU/L (reference range 4–51 IU/L) and total bilirubin of 31 $\mu\text{mol/L}$ (0–21 $\mu\text{mol/L}$). A summary trend of his liver markers is presented in figure 1. An extended liver panel, including hepatitis A, B, C and E, syphilis, autoimmune antibodies, ferritin, alpha-1 antitrypsin and caeruloplasmin

screen, was negative or normal. HIV-1 viral load remained undetectable (under 20 copies/mL) throughout.

He received a 7-day course of doxycycline, as well as metronidazole and clindamycin for an asymptomatic chlamydia infection and a perianal abscess during this period, respectively; however, both diagnoses and treatments were given after the increase in liver markers. He reported no regular prescription, over-the-counter or herbal medications, and no increased alcohol intake. Liver ultrasound showed an echogenic liver in keeping with mild fatty infiltration. On liver transient elastography, the median liver stiffness was 5.4 kPa and the controlled attenuation parameter was 205, suggesting no significant fibrosis or steatosis.

Histology revealed diffuse confluent centrilobular necrosis, which is a typical lesion of drug-induced liver injury but not specific, present also in new-onset autoimmune hepatitis or in some cases of hepatitis virus infection, the two main differential diagnoses. Both conditions were reasonably excluded in our patient.

The patient remained asymptomatic apart from mild nausea after antiretrovirals. Rilpivirine was switched to raltegravir 1200 mg once daily on week 66. His ALT showed an improvement soon after the switch; however, raltegravir was switched to boosted darunavir with ritonavir on week 71 due to poor mood. Abacavir/lamivudine was subsequently switched to tenofovir disoproxil fumarate/emtricitabine on week 74 due to nausea. By week 92, liver markers had normalised and his nausea resolved.

Low rates of liver toxicity have been reported with rilpivirine. In the ECHO and THRIVE studies comparing rilpivirine

with efavirenz, rilpivirine had significantly lower incidence of grade 2–4 ALT and aspartate transaminase levels,³ with no serious treatment-related hepatic adverse events. One other case of possible drug-induced allergic hepatitis secondary to rilpivirine has been reported⁴; however, the patient was severely immunosuppressed with disseminated *Mycobacterium avium* infection, multiple comorbidities and concomitant medications, without histological evidence of underlying liver pathology. While rilpivirine-induced liver damage is rare, further monitoring and reporting of such events are important to better understand the safety profile of this drug.

Ming Jie Lee ¹, Philip Berry,² Francesca D'Errico,² Rosa Miquel,³ Ranjababu Kulasegaram¹

¹Harrison Wing Department, Guy's and St Thomas Hospital NHS Foundation Trust, London, UK

²Department of Gastroenterology and Hepatology, Guy's and St Thomas Hospital NHS Foundation Trust, London, UK

³Liver Histopathology Laboratory, Institute of Liver Studies, King's College Hospital NHS Foundation Trust, London, UK

Correspondence to Dr Ming Jie Lee, Harrison Wing Department, Guy's and St Thomas Hospital NHS Foundation Trust, London SE1 9RS, UK; minglee@doctors.org.uk

Handling editor Anna Maria Geretti

Twitter Ming Jie Lee @Jnim87

Acknowledgements The authors would like to acknowledge the person in the case for his consent and contribution to the case report.

Contributors MJL and RK contributed to the draft and final review of the article. PB, FDE and RM contributed to the review and editing of the article and to its resubmission.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Obtained.

Provenance and peer review Not commissioned; internally peer reviewed.

© Author(s) (or their employer(s)) 2020. No commercial re-use. See rights and permissions. Published by BMJ.

To cite Lee MJ, Berry P, D'Errico F, et al. *Sex Transm Infect* 2020;**96**:618–619.

Received 24 October 2019
Revised 17 December 2019
Accepted 12 January 2020
Published Online First 23 January 2020

Sex Transm Infect 2020;**96**:618–619.
doi:10.1136/sextrans-2019-054336

ORCID iD

Ming Jie Lee <http://orcid.org/0000-0002-5244-0070>

Figure 1 Liver markers and association with antiretroviral switches over time. ABC/3TC, abacavir and lamivudine fixed-dose combination; ALT, alanine transaminase; BD, twice daily; Bil, total bilirubin; DRV/r, boosted darunavir with ritonavir; IU, international unit; OD, once daily; RAL, raltegravir; RPV, rilpivirine; TDF/FTC, tenofovir disoproxil fumarate and emtricitabine fixed-dose combination; ULN, upper limit of normal range.

REFERENCES

- 1 Weber R, Sabin CA, Friis-Møller N, *et al.* Liver-related deaths in persons infected with the human immunodeficiency virus: the D:A:D study. *Arch Intern Med* 2006;166:1632–41.
- 2 Núñez M. Hepatotoxicity of antiretrovirals: incidence, mechanisms and management. *J Hepatol* 2006;44:S132–9.
- 3 Nelson M, Amaya G, Clumeck N, *et al.* Efficacy and safety of rilpivirine in treatment-naïve, HIV-1-infected patients with hepatitis B virus/hepatitis C virus coinfection enrolled in the phase III randomized, double-blind echo and thrive trials. *J Antimicrob Chemother* 2012;67:2020–8.
- 4 Ahmed Y, Siddiqui W, Enoch CB, *et al.* Rare case of rilpivirine-induced severe allergic hepatitis. *J Antimicrob Chemother* 2013;68:484–6.