

Research letter

HIV, HBV and syphilis screening in antenatal care in Lubango, Angola

Screening for infections is part of antenatal care (ANC) evidence-based interventions.¹ The Angola's Health Development Plan 2012–2025² includes screening of human immunodeficiency virus (HIV), hepatitis B virus (HBV) and syphilis in the essential package of interventions for maternal and newborn health, which is done using an opt-out approach. There are only a few reports about ANC in Angola.^{3–5} To our knowledge, this is the first one about ANC infection screening. We characterised ANC screening rates for HIV, HBV and syphilis among 500 parturient women admitted between October 2016 to September 2017 to the Irene Neto Maternity service in Lubango, Huíla, the second most populous province of Angola (Oliveira *et al*, 2020, accepted to *Sex Transm Infect*). One additional aim was to explore rural versus urban disparities in screening rates, according to parturients place of residence, and identify opportunities for improvement. Since no demographic surveillance system was available to us, rural or urban residence was determined by parturient description of her place of residence. Every parturient residing in Lubango or in its suburbs was considered to live in an urban area. On the contrary, those living outside this perimeter, whose source of livelihood was

mostly agriculture, were considered to be from a rural area. Whenever they could not read or write, pregnant woman's fingerprint and witness signature were required, after informed consent was read to her (Oliveira *et al*, 2020, accepted to *Sex Transm Infect*). Timing and number of ANC visits and results of ANC infection screening were retrieved from each participant's pregnancy card. When a positive screening result was found, the woman was asked about the treatment received. At admission in labour, 6.6% (33/500) of the women did not present a pregnancy card. Of the remaining 467 women, some presented pregnancy cards that were only partially legible (reflected in the denominators below). One way of assessing the quality of ANC is to estimate the proportion of pregnant women initiating ANC during the first trimester of pregnancy.³ Overall, 9.1% (42/462) of the women had initiated ANC in the first trimester of pregnancy (table 1), without difference based on rural or urban residence. In 2009, WHO recommended a minimum of four ANC visits and, in 2017, updated to a minimum of eight contacts to reduce perinatal mortality and improve women's experience of care.^{6 7} The Angola's Health Development Plan 2012–2025 recommends that a pregnant woman should be observed in, at least, four ANC visits.² In this study, at least four ANC visits during pregnancy were observed in 62.7% (293/467). HIV, HBV and syphilis screening was performed in 66.5% (309/465), 44.6% (208/466) and 60.2% (281/467) respectively, and all infections screening rates were

significantly lower in women from rural areas compared with those from urban areas. Only 2.6% (12/465), 0.6% (3/466) and 0.6% (3/467) of the parturients were tested two times during pregnancy against HIV, HBV and syphilis, respectively. Of the five women who had tested HIV positive, one referred that had not initiated antiretroviral therapy. Penicillin G treatment was reported by five of the seven women who tested positive for syphilis. Among the five treated with penicillin G, three recalled that had received the recommended three doses. None of those who tested positive for HBV referred having initiated treatment. In this study, the timing and number of ANC visits fell below the expectations set by the WHO^{6 7} as also, although not so low, found in previous studies conducted in other Angola provinces.^{3 5} In line with findings of previous studies,^{8–10} rural women presented significantly lower rates of HIV, HBV and syphilis screening. This could be explained by the fact that in Angolan rural areas, the offer of health services in general is lower, particularly laboratory tests. Additionally, women from rural areas have a lower educational level, which probably makes them less able to communicate with and access healthcare services. Policies are required to improve the quality of ANC in Lubango and to address rural–urban disparities in ANC infection screening rates.

Dinamene Oliveira^{1,2},
 Maria do Rosário Martins,² Rita Castro,²
 Lemuel Cordeiro,³ Maria Rosalina Barroso,⁴
 Maria Antónia Nazaré,⁵ Filomena Pereira²

Table 1 ANC characteristics: general and according with area of residence

| ANC characteristics | Total | | Area of residence | | P value |
|--|------------|--------------|-------------------|-------------|---------|
| | N (%) | 95% CI | Urban n (%) | Rural n (%) | |
| Timing and number of visits | | | | | |
| First trimester (n=462) | 42 (9.1) | 6.8 to 12.1 | 35 (8.7) | 7 (11.9) | 0.428* |
| ≥4 visits (n=467) | 293 (62.7) | 58.3 to 67.0 | 262 (64.2) | 31 (52.5) | 0.083* |
| ≥8 visits (n=467) | 16 (3.4) | 2.1 to 5.5 | 14 (3.4) | 2 (3.4) | 1.000† |
| Infection screening documented (at least once) | | | | | |
| HIV (n=465) | 309 (66.5) | 62.0 to 70.6 | 285 (70.2) | 24 (40.7) | <0.001* |
| HBsAg (n=466) | 208 (44.6) | 40.2 to 49.2 | 201 (49.4) | 7 (11.9) | <0.001* |
| RPR (n=467) | 281 (60.2) | 55.7 to 64.5 | 269 (65.9) | 12 (20.3) | <0.001* |
| Infection treatment reported by women | | | | | |
| HIV positive (n=5) | 4 (80.0) | 37.6 to 96.4 | 4 (80.0) | 0 (0.0) | NA |
| HBsAg positive (n=10) | 0 (0.0) | 0.0 to 27.8 | NA | NA | NA |
| RPR positive (n=7) | 5 (71.4) | 35.9 to 91.8 | 4 (80.0) | 1 (50) | NA |

*Pearson χ^2 test.

†Fisher's exact test.

ANC, antenatal care; n, absolute frequencies; %, relative frequencies; 95% CI, 95% confidence intervals (estimated with the Wilson method using EpiTools); HBsAg, hepatitis B surface antigen; RPR, rapid plasma reagin test; NA, not applicable.

¹Posto Médico do Lubango, Clínica Girassol, Lubango, Huíla, Angola
²Global Health and Tropical Medicine, Universidade Nova de Lisboa Instituto de Higiene e Medicina Tropical, Lisboa, Portugal
³Gabinete de Ensino, Pesquisa e Pós-graduação, Clínica Girassol, Luanda, Angola
⁴Unidade de Cuidados Intensivos Neonatais, Hospital Prof. Doutor Fernando Fonseca, Lisboa, Portugal
⁵Departamento da Mulher, Hospital Prof. Doutor Fernando Fonseca, Lisboa, Portugal

Correspondence to Dinamene Oliveira, Posto Médico do Lubango, Clínica Girassol, Lubango, Huíla, Angola; odinamene@gmail.com

Handling editor Anna Maria Geretti

Acknowledgements The authors are grateful to the parturients that participated in the study. We also want to thank the staff of the Irene Neto Maternity, particularly the nurses team.

Contributors Conceived and designed the study: all authors. Performed the data collection: DO. Performed the screening tests: DO and RC. Analysed the data: DO and MdRM. Wrote the paper: DO, FP and MdRM. Revised the paper: RC, MRB, MAN and LC.

Funding Calouste Gulbenkian Foundation supported this study through a scholarship awarded to the corresponding author (grant number 135499).

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval Parturients had to sign an informed consent form to be included in this study, which was approved by the Angola Ethics Committee (reference 35/2017).

Provenance and peer review Not commissioned; internally peer reviewed.

© Author(s) (or their employer(s)) 2020. No commercial re-use. See rights and permissions. Published by BMJ.



To cite Oliveira D, Martins MdR, Castro R, *et al.* *Sex Transm Infect* 2020;**96**:621–622.

Received 18 December 2019

Revised 4 March 2020

Accepted 18 March 2020

Published Online First 31 March 2020

Sex Transm Infect 2020;**96**:621–622.

doi:10.1136/sextrans-2019-054407

ORCID iD

Dinamene Oliveira <http://orcid.org/0000-0002-7765-5991>

REFERENCES

1 The Partnership for Maternal, Newborn & Child Health (PMNCH). *A global review of the key interventions related to reproductive, maternal, newborn and child health (RMNCH)*. Geneva: PMNCH, 2011.

- Ministério da Saúde (MINSa). *Plano Nacional de Desenvolvimento Sanitário 2012-2025*. Luanda: MINSa, 2012.
- Nimi T, Fraga S, Costa D, *et al.* Prenatal care and pregnancy outcomes: a cross-sectional study in Luanda, Angola. *Int J Gynaecol Obstet* 2016;135:572–8.
- Instituto Nacional de Estatística (INE), Ministério da Saúde (MINSa), Ministério do Planeamento e do Desenvolvimento Territorial (MINPLAN), ICF. *Inquérito de Indicadores Múltiplos e de Saúde em Angola 2015-2016*. Luanda and Rockville: INE, MINSa, MINPLAN and ICF, 2017.
- Rosário EVN, Gomes MC, Brito M, *et al.* Determinants of maternal health care and birth outcome in the Dande health and demographic surveillance system area, Angola. *PLoS One* 2019;14:e0221280.
- World Health Organization. *WHO Recommended Interventions for Improving Maternal and Newborn Health*. Geneva: WHO, 2009.
- World Health Organization. *WHO recommendations on maternal health: guidelines Approved by the who guidelines review Committee*. Geneva: WHO, 2017.
- Ousman SK, Mdala I, Thorsen VC, *et al.* Social determinants of antenatal care service use in Ethiopia: changes over a 15-year span. *Front Public Health* 2019;7:161.
- Yaya S, Bishwajit G, Shah V. Wealth, education and urban-rural inequality and maternal healthcare service usage in Malawi. *BMJ Glob Health* 2016;1:e000085.
- Tran TK, Nguyen CTK, Nguyen HD, *et al.* Urban - rural disparities in antenatal care utilization: a study of two cohorts of pregnant women in Vietnam. *BMC Health Serv Res* 2011;11:120.