

# Factors associated with lower knowledge of HIV and STI transmission, testing and treatment among MSM in Ireland: findings from the MSM Internet Survey Ireland (MISI) 2015

Cian Carey <sup>1,2</sup>, Kate O'Donnell,<sup>3</sup> Martin Davoren,<sup>2,4</sup> Mick Quinlan,<sup>5</sup> Derval Igoe,<sup>3</sup> Peter Barrett<sup>2,6</sup>

► Additional material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/sextrans-2020-054469>).

<sup>1</sup>School of Medicine, University College Cork, Cork, Ireland

<sup>2</sup>School of Public Health, University College Cork, Cork, Ireland

<sup>3</sup>HSE Health Protection Surveillance Centre, Dublin, Ireland

<sup>4</sup>Sexual Health Centre, University College Cork, Cork, Ireland

<sup>5</sup>Gay Health Network, Dublin, Ireland

<sup>6</sup>Department of Public Health HSE South, St. Finbarr's Hospital, Cork, Ireland

## Correspondence to

Dr Peter Barrett, School of Public Health, University College Cork College of Medicine and Health, Cork, Ireland; [peterbarrett1@hotmail.com](mailto:peterbarrett1@hotmail.com)

Received 4 February 2020

Revised 1 September 2020

Accepted 3 October 2020

Published Online First

26 October 2020



© Author(s) (or their employer(s)) 2021. No commercial re-use. See rights and permissions. Published by BMJ.

**To cite:** Carey C, O'Donnell K, Davoren M, et al. *Sex Transm Infect* 2021;**97**:351–356.

## ABSTRACT

**Background** Poor knowledge regarding STI and HIV transmission, testing and treatment among men who have sex with men (MSM) may be contributing to their disproportionate burden of STIs. However, factors which predispose MSM to having this low knowledge are less understood.

**Aim** The aim of this study was to identify factors associated with having lower knowledge of HIV and STI transmission, testing and treatment among MSM.

**Methods** The MSM Internet Survey Ireland 2015 was a self-completed online national survey available to MSM living in Ireland. Thirteen factual statements were used to assess participants' knowledge of HIV and STI transmission, testing and treatment. Respondents were defined as having 'lower knowledge' if they indicated prior knowledge of fewer than 11 true statements. Multivariable-adjusted logistic regression was used to identify factors associated with having lower knowledge.

**Results** 3090 MSM completed the survey, of whom 2905 (94%) were included in this study. Thirty-six per cent (n=1055) had lower knowledge of HIV and STI transmission, testing and treatment. The factors associated with having lower knowledge included being 18–24 years of age (vs >40 years; adjusted OR (aOR) 1.98, 95% CI 1.50 to 2.61), born in Ireland (vs outside Ireland; aOR 1.62, 95% CI 1.25 to 2.10) and being out to none of their contacts (vs out to most/all; aOR 1.69, 95% CI 1.20 to 2.37). Knowledge was also lower among those who never tested for HIV (vs tested negative within last 12 months; aOR 2.32, 95% CI 1.88 to 2.86) and among MSM who never visited an MSM-specific health promotion website (vs visited website; aOR 1.81, 95% CI 1.45 to 2.25).

**Conclusion** A range of demographic factors, sexual health variables and use of MSM-specific sexual health promotion materials are associated with low knowledge regarding HIV and STI transmission, testing and treatment among MSM in Ireland. Engagement with the main national MSM-specific sexual health promotion website was associated with higher knowledge levels.

## BACKGROUND

STIs disproportionately affect men who have sex with men (MSM).<sup>1</sup> Although MSM only represent

approximately 6% of the Irish male population, they accounted for 49% of HIV, 86% of early infectious syphilis and 64% of gonorrhoea diagnoses in the overall population in Ireland in 2018.<sup>2,3</sup> This follows a similar trend throughout Europe and in the USA, where in recent years MSM have accounted for 50% and 67% of newly diagnosed HIV cases, respectively.<sup>4,5</sup>

Previous research highlights that MSM are more likely to experience an earlier sexual debut, are more likely to have multiple concurrent sexual partners and report longer cumulative lifetime periods of new partner acquisition when compared with heterosexual men.<sup>6,7</sup> These behavioural issues, coupled with the fact that the risk of HIV transmission from condomless anal intercourse (CAI) is higher than unprotected vaginal sex, partially explain the higher rates of HIV in this group.<sup>8</sup> In addition, many sociostructural issues, including the stigmatisation of the MSM population, have been shown to increase MSM vulnerability for HIV acquisition.<sup>9,10</sup>

Improved knowledge of HIV and STI transmission pathways is associated with increased awareness of preventive interventions and reduced spread of HIV/STI among sexual partners.<sup>11</sup> Although knowledge is only one component of health literacy, previous research has highlighted the positive association between HIV-related knowledge and overall health literacy levels.<sup>12</sup> It remains relevant and timely to conduct dedicated research on knowledge of HIV/STI among MSM, given the advances made in prevention and treatment in recent years. Younger MSM are growing up in a post-AIDS era, with ever-increasing availability of both treatment as prevention and pre-exposure prophylaxis (PrEP) for HIV, but with a growing threat from treatment-resistant STIs. Public health interventions, which can address low levels of HIV and STI-related knowledge at an individual level, have the potential to confer considerable benefits. A previous systematic review reported that MSM who received online educational interventions reported improved knowledge levels and had lower incidence of CAI with non-steady partners subsequently.<sup>13</sup>

Few studies from the European context have been able to clearly identify which subgroups of MSM

have lower knowledge levels, and existing studies have been limited by relatively small sample sizes or by assessing participants' knowledge of HIV alone, without considering other STIs.<sup>14 15</sup> The only previous study in an Irish context focused on MSM and heterosexuals together, and while it highlighted knowledge deficits regarding HIV transmission, there were too few MSM respondents for a dedicated analysis.<sup>16</sup>

Using data from the MSM Internet Survey Ireland (MISI 2015), the aims of this study were to compare MSM who had lower levels of knowledge of HIV and STI transmission, testing and treatment with MSM who did not, and to identify factors which may predispose them to having lower knowledge levels.

## METHODS

### Study survey design

We used data from the MISI 2015, which were an anonymous, cross-sectional survey of MSM living in Ireland. The survey was accessible online from 1 March to 31 May 2015 and was conducted prior to the introduction of PrEP in Ireland in 2017. The survey, which included 112 questions, was only available in English and took approximately 14 min to complete. Questions in MISI 2015 were derived from previous surveys, including the UK Gay Men's Sex Survey 2014 and the European MSM Internet Survey 2010. Certain questions were adapted and added to make the survey relevant to an Irish setting. Further details regarding the survey methodology are available in the MISI 2015 survey report.<sup>17</sup>

### Study population and sample

The inclusion criteria for MISI 2015 were as follows: living in Ireland, identifying as a man or trans man, 18 years or older, sexually attracted to men and/or previously had sex with men or expected to in the future.

Survey participants were recruited via advertising on lesbian, gay, bisexual and transgender (LGBT+), STI and health promotion websites, through social media campaigns and through promotional card distribution at gay social and community venues. A launch party was held to initially promote the survey, with subsequent advertisements placed in national media and banner advertisements and emails used to target subscribers on gay networking apps. The majority (61%) of men accessed the survey via the MISI website directly, with the breakdown of visitor numbers from recruiting websites available in the full survey report.<sup>17</sup> Further detailed information on the survey promotion has been described elsewhere.<sup>18 19</sup>

### Survey measurement

Lower knowledge of HIV and STI transmission, testing and treatment was our primary outcome of interest. Knowledge was assessed in this survey using 13 factually correct statements, which are presented in box 1. These statements were used to examine men's overall HIV literacy and were not used as an examination of their personal HIV risk reduction strategies. The first seven statements related to an individual's prior knowledge of HIV testing and treatment, while the other six examined an individual's knowledge of transmission of HIV and STIs. After reading each statement, participants were asked to specify their prior knowledge by selecting one of five responses: *I knew this already/I was not sure about this/I did not know this already/I do not understand this/I do not believe this* (online supplemental table 1).

Using participants' responses for each of these statements, we derived a binary variable for each statement of 'knew' (*I knew*

### Box 1 Statements used to assess men who have sex with men knowledge of HIV/STI transmission, testing and treatment

Statements regarding knowledge of HIV, testing and treatment (used in primary analysis only)

1. AIDS is caused by a virus called HIV.
2. You cannot be confident about whether someone has HIV or not from their appearance.
3. There is a medical test that can show whether or not you have HIV.
4. If someone becomes infected with HIV, it may take several weeks before it can be detected in a test.
5. There is currently no cure for HIV infection.
6. HIV infection can be controlled with medicines so that its impact on health is much less.
7. Effective treatment of HIV infection reduces the risk of HIV being transmitted.

Statements regarding HIV/STI transmission (used in primary and secondary analyses)

1. HIV cannot be passed during kissing, including deep kissing, because saliva does not transmit HIV.
2. People can pick up HIV through their penis while being 'active' in unprotected intercourse (fucking) with an infected partner even if they don't ejaculate.
3. People can pick up HIV through their rectum while being passive in unprotected intercourse (being fucked) with an infected partner.
4. Even without ejaculation, oral sex (sucking and being sucked) carries a risk of infection with syphilis or gonorrhoea.
5. When HIV infected and uninfected people have sex together, the chances of HIV being passed on are greater if either partner has another STI.
6. Most STIs can be passed on more easily than HIV.

*this already* response) or 'did not know' (any of the other four responses). We next calculated the total number of statements each participant knew out of 13. Finally, we developed a binary variable of 'lower knowledge' if the respondent knew 10 statements or fewer out of 13. Eleven or more out of 13 was defined as 'higher knowledge'. As MISI 2015 was a new survey, this cut-off was chosen to ensure an adequate number of respondents in each group and to reduce the risk of a type II error. This binary lower knowledge variable was then included as the dependent variable in the primary analysis. A dichotomous outcome variable was chosen (lower vs higher knowledge) for analysis due to skewed distribution in the numbers of men who knew each number of statements (online supplemental table 2). The use of a continuous outcome would have required a log (or other) transformation to be applied, and this may have impacted on the ability for clear public health messages to be drawn from the results.

We undertook a secondary analysis which focused on participants' knowledge of HIV/STI transmission (excluding statements related to testing or treatment). A previous cross-sectional study demonstrated poor knowledge of HIV/STI transmission among the general Irish population, and thus, this warranted inclusion as a separate analysis.<sup>16</sup> This was performed using the six HIV and STI transmission statements that were included among the original 13 statements as discussed previously (last 6 statements in box 1). We again derived a new binary variable for lower versus higher knowledge, where lower knowledge

for this analysis was four or fewer out of six, with five or more correct answers deemed to indicate higher knowledge of HIV/STI transmission.

### Data analysis

Primary analysis was restricted to respondents who answered all 13 questions about HIV/STI transmission, testing and treatment.

Associations between a range of demographic factors, exposure to health promotional materials and other variables pertaining to sexual health and orientation with overall knowledge levels were investigated. These covariates were selected a priori based on previous studies, which suggested associations with overall knowledge of HIV and risky sexual behaviours.<sup>19 20</sup>

The included demographic and sexual health/orientation factors can be seen in [box 1](#) and are described in the supplement.

Furthermore, we included two binary covariates in the models relating to men's exposure to specific health promotion materials. These materials included the national government-supported MSM-specific sexual health promotion website ([www.man2man.ie](http://www.man2man.ie)) and four infographics, which had been previously developed by community partners to provide sexual health advice for MSM in Ireland (online supplemental file 1). The infographics covariate was derived as having seen at least one of the four infographics or having seen none of them previously. Similarly, we included a covariate for having visited the MSM-specific website or not.

Univariable analysis was performed between each of these covariates and lower knowledge among MSM. ORs were calculated with 95% CIs and two-sided significance set at  $p < 0.05$ . We considered the overall proportion of MSM with lower knowledge to be the most relevant outcome, rather than count data, and thus, we used logistic regression rather than log-linear models (eg, Poisson regression). Subsequently, multivariable-adjusted logistic regression analysis was undertaken, with all covariates included in adjusted models. This selection was based on a consensus decision-making process with health service and community partners, and informed by previous literature.<sup>15</sup> Adjusted ORs (aORs) with 95% CIs were calculated to identify independent covariates that remained associated with lower knowledge. Where a variable had  $>5\%$  missing data, a missing indicator covariate was created and included in the analyses.

For our secondary analysis, we assessed participants' knowledge of HIV/STI transmission alone with the same factors described earlier. The results of this are provided in the supplement (including online supplemental table 3).

A sensitivity analysis assessing low knowledge among only HIV-negative MSM was also conducted, with the results available in online supplemental table 4.

In addition, collinearity between independent covariates was tested. In the correlation matrices of predictor covariates, all pairwise Pearson correlation coefficients were  $<0.4$ , suggesting that collinearity was not an important factor in this study. All data analyses were undertaken in StataSE V.15.

### RESULTS

The number of valid responses to MISI 2015 was 3090. We excluded 185 men who did not answer all 13 questions relating to HIV/STI treatment, transmission and testing, and our final sample was 2905 men (94% of total). The median age was 30 years (range 18–78 years), with 5% ( $n=142$ ) HIV positive. Ninety per cent ( $n=2611$ ) of men had sex with another man within the previous 12 months. The median knowledge score for study participants was 11 out of 13 (IQR 10–13) (online

supplemental table 2). A comparison between our study sample and the excluded men is shown in online supplemental table 5.

### Lower knowledge

One thousand and fifty-five men (36%) had lower knowledge of HIV/STI testing, transmission and treatment. In univariable analysis, factors associated with having lower knowledge are shown in [table 1](#).

From the multivariable analysis, lower knowledge was significantly more likely among men aged 18–24 years (aOR 1.98, 95% CI 1.50 to 2.61), those educated below degree level (aOR 1.58, 95% CI 1.32 to 1.88), men living outside Dublin (aOR 1.21, 95% CI 1.01 to 1.44), those born in Ireland (aOR 1.62, 95% CI 1.25 to 2.10) and among men who were unsure of their HIV status (aOR 1.35, 95% CI 1.13 to 1.61). Lower levels of knowledge were also observed among men who were out to fewer people, with significantly higher aORs observed in those out to few (aOR 1.36, 95% CI 1.06 to 1.76) or none (aOR 1.69, 95% CI 1.20 to 2.37) compared with those out to all/almost all. MSM who had never tested for HIV also had lower knowledge levels compared with those who last tested negative for HIV within the previous 12 months (aOR 2.32, 95% CI 1.88 to 2.86). Furthermore, non-engagement with the national MSM-specific health promotion website was associated with lower knowledge (aOR 1.81, 95% CI 1.45 to 2.25). Full results, including the baseline comparison group for each covariate, can be seen in [table 1](#).

Of note from the secondary analysis, in multivariable-adjusted models, students had low knowledge levels of HIV/STI transmission compared with men who were employed (aOR 1.35, 95% CI 1.07 to 1.71). In addition, age, education level, country of birth, outness level, HIV testing history, certainty of HIV status and prior non-engagement with the national MSM-specific health promotion website were significantly associated with knowledge levels. In contrast to the primary analysis, men living outside of Dublin did not appear to have low knowledge of HIV transmission compared with men living in Dublin (aOR 1.04, 95% CI 0.88 to 1.23). The full results of the secondary analysis can be seen in the supplemental results section (online supplemental table 3).

### DISCUSSION

#### Principal findings

Our primary analysis included 2905 MSM living in Ireland in 2015, before the widespread availability of PrEP. Although the majority of MSM in Ireland had high levels of knowledge about HIV/STI transmission, testing and treatment, a large minority (36%) had low knowledge. While knowledge is only a single component of health literacy, its importance has been documented in prior studies which have shown that knowledge of HIV is a good predictor of a person's overall health literacy.<sup>12</sup> Previous research has highlighted the benefits of interventions aimed at improving knowledge of HIV/STI transmission, testing and treatment among MSM, with some evidence of subsequent reductions in risky sexual behaviour.<sup>13</sup>

In total, 56% of MSM who had never tested for HIV had lower knowledge compared with only 9% of HIV-positive MSM. These results are similar to those from existing studies which reported that HIV-positive MSM tend to have better knowledge about HIV treatment and onward transmission risks, likely due to their HIV diagnosis acting as an educational intervention.<sup>14</sup> A similar trend was observed in our secondary analysis, which focused on HIV/STI transmission only. Lack of knowledge about HIV

## Behaviour

**Table 1** Univariable and multivariable analyses of lower knowledge of HIV and STI transmission, testing and treatment among MSM in Ireland, MISI 2015

	Number with lower knowledge (%)	Univariable analysis			Multivariable analysis		
		OR	95% CI	P value	aOR	95% CI	P value
Age group (years)							
>40	234 (30)	1	Ref	—	1	Ref	—
25–39	353 (29)	0.98	(0.80 to 1.19)	0.82	1.22	(0.98 to 1.53)	0.080
18–24	468 (51)	2.48	(2.03 to 3.02)	<0.001	1.98	(1.50 to 2.61)	<0.001
Education							
Degree or higher	431 (28)	1	Ref	—	1	Ref	—
Less than a degree	601 (46)	2.20	(1.88 to 2.57)	<0.001	1.58	(1.32 to 1.88)	<0.001
Area of residence*							
Dublin	403 (30)	1	Ref	—	1	Ref	—
Outside Dublin	589 (42)	1.69	(1.44 to 1.97)	<0.001	1.21	(1.01 to 1.44)	0.034
Missing	63 (34)	1.20	(0.86 to 1.66)	0.281	1.05	(0.74 to 1.51)	0.771
Employment							
Employed	580 (31)	1	Ref	—	1	Ref	—
Unemployed	83 (44)	1.74	(1.29 to 2.36)	<0.001	1.23	(0.87 to 1.72)	0.237
Student	322 (49)	2.18	(1.82 to 2.62)	<0.001	1.10	(0.86 to 1.41)	0.452
Other†	49 (40)	1.51	(1.04 to 2.19)	0.032	1.55	(1.02 to 2.35)	0.039
Country of birth							
Outside Ireland	106 (26)	1	Ref	—	1	Ref	—
Ireland	944 (38)	1.78	(1.41 to 2.25)	<0.001	1.62	(1.25 to 2.10)	<0.001
Sexual Identity							
Gay	756 (33)	1	Ref	—	1	Ref	—
Bisexual	185 (48)	1.81	(1.45 to 2.24)	<0.001	1.04	(0.80 to 1.37)	0.756
Straight/other	98 (45)	1.64	(1.24 to 2.17)	0.001	0.90	(0.65 to 1.25)	0.523
Outness level							
Out to all/almost all	433 (30)	1	Ref	—	1	Ref	—
More than half	155 (37)	1.35	(1.08 to 1.70)	0.009	1.16	(0.90 to 1.48)	0.256
Less than half	94 (36)	1.32	(1.00 to 1.75)	0.047	1.03	(0.76 to 1.41)	0.832
Few	206 (44)	1.86	(1.50 to 2.31)	<0.001	1.36	(1.06 to 1.76)	0.017
None	136 (54)	2.70	(2.06 to 3.54)	<0.001	1.69	(1.20 to 2.37)	0.002
HIV testing history							
Negative <12 months	267 (25)	1	Ref	—	1	Ref	—
Negative >12 months	169 (29)	1.23	(0.98 to 1.54)	0.074	1.14	(0.89 to 1.46)	0.285
Positive	13 (9)	0.31	(0.17 to 0.56)	<0.001	0.38	(0.21 to 0.69)	0.001
Never tested	598 (56)	3.96	(3.30 to 4.76)	<0.001	2.32	(1.88 to 2.86)	<0.001
STI in the last 12 months							
No/don't know	986 (38)	1	Ref	—	1	Ref	—
Yes	62 (23)	0.48	(0.36 to 0.65)	<0.001	0.89	(0.64 to 1.23)	0.486
Certainty of HIV status							
Sure of HIV status	648 (33)	1	Ref	—	1	Ref	—
Unsure of HIV status	400 (43)	1.50	(1.28 to 1.76)	<0.001	1.35	(1.13 to 1.61)	0.001
Visited Man2Man website				<0.001	<0.001		
Visited	178 (24)	1	Ref	—	1	Ref	—
Not visited	855 (41)	2.24	(1.85 to 2.70)	<0.001	1.81	(1.45 to 2.25)	<0.001
Viewed one or more HP infographics				<0.001	0.022		
Viewed	465 (32)	1	Ref	—	1	Ref	—
Not viewed	551 (40)	1.46	(1.25 to 1.71)	<0.001	1.17	(0.98 to 1.41)	0.084

\*Variable with &gt;5% missing data.

†Other includes retired, sick leave, medically retired and other.

aOR, adjusted OR; HP, health promotion; MISI, MSM Internet Survey Ireland; MSM, men who have sex with men; Ref, reference.

transmission is associated with irregular condom use and higher numbers of sexual partners.<sup>15</sup> This may contribute to potential undiagnosed HIV and STIs among this cohort of MSM. It is notable that over 25% of men who previously tested negative for HIV also had low knowledge levels. This is concerning as it

appears that the opportunity for educating these men about HIV and STIs at the time of testing is being missed. It has been shown that MSM are likely to practice safer sex in the immediate aftermath of HIV testing and so perhaps may also be more receptive to long-term knowledge improving interventions at this time.<sup>21</sup>



Over one-third (37%) of the sample had never tested for HIV and 32% disclosed that they were unsure of their HIV status. The proportion of men who had never tested for HIV was higher than reported in other recent surveys of MSM, and it may be partially explained by the relatively high response rates from younger MSM in MISI 2015.<sup>19</sup> Based on the potential burden of undiagnosed HIV and STIs, non-testers and infrequent HIV testers are an important group to target for knowledge improvement. This requires interventions which increase awareness about the relevance of testing and which improve access to testing as a priority. A previous study highlighted the benefit of community-based peer testing services for hard-to-reach groups like MSM, and similar interventions in an Irish setting may improve testing uptake among MSM.<sup>22</sup>

The lower levels of knowledge of HIV/STI transmission (from our secondary analysis) among students is a cause for concern. Heavy alcohol consumption in this group has been linked to inconsistent condom use previously.<sup>23,24</sup> It is plausible that lower levels of knowledge of HIV/STI transmission among students may also be contributing to high levels of risky sexual behaviour and onward spread of HIV and STIs.<sup>25</sup> Improved knowledge about HIV transmission risk may improve condom use among students.<sup>26</sup> The knowledge disparity among younger MSM also underlines the importance of providing interventions to educate younger MSM, and all students, on safer sexual practices in schools and third-level institutions. Multiple frameworks for how to develop and deliver such teaching have been produced previously and may be adapted to the Irish setting.<sup>27</sup> These frameworks provide information for educators on how to maximise student participation in educational programmes and are associated with improved knowledge among young people.

MSM who disclosed their sexuality to few or no people had lower knowledge of HIV/STI transmission, testing and treatment when compared with MSM who were openly out to all or almost all. MSM who are out to more people are also more likely to disclose this fact to their medical practitioners and are more likely to seek information on topics relating to sexual health.<sup>28</sup> While interventions could be tailored to MSM who are less likely to engage with sexual health services, interventions which aid MSM being more open about their sexual orientation are preferable. Stigma around LGBT+ issues still exists in Irish society and may negatively affect men's confidence in disclosing their sexual orientation.<sup>29</sup> It is thus important that emphasis is placed on support that is known to help improve 'outness', including mental health support, through the creation of LGBT+ social spaces. These interventions, in addition to the implementation of a National LGBT+ Inclusion Action Plan, may help accelerate acceptance of LGBT+ issues nationwide and increase individuals' willingness to be open about their sexual orientation.

One potential option for reaching MSM is by making general health promotion materials accessible and appealing to larger audiences. Our findings show that a national health promotion website can be effective, perhaps more so than infographics or information leaflets in engaging MSM. Widespread promotion of health promotion materials, especially websites, may be beneficial for knowledge improvement, as research has shown that MSM may benefit from similar online interventions.<sup>13</sup> However, it is important to consider the difficulties of evaluating the effectiveness of these materials and the potential limitations of selection bias among those who are likely to engage with such materials.

## Study limitations

While MISI 2015 is currently the largest ever study of the MSM population in Ireland, it is not without limitations. First, we regret that we were unable to include a risk behaviour covariate (eg, number of sexual partners). However, this was unavoidable due to an intrinsic limitation in the way that data on the number of sexual partners were collected in MISI 2015. The learning from this survey has been used to inform the design of subsequent behavioural surveys of MSM in Ireland. In addition, all data were self-reported, with the survey only available online and through English. A convenience sample was used to maximise the size of the recruited sample. This may have introduced selection bias into the study, especially biasing the sample towards those who had computer access. However, previous research has highlighted that online surveys are the optimal method for recruiting a generalisable MSM sample, and so online recruitment is unlikely to have had major impact on the representativeness of our study population.<sup>30</sup> Furthermore, we excluded participants who did not answer all 13 statements pertaining to HIV/STI knowledge. However, we compared the demographics of excluded participants with our study sample and found no major differences between the groups. We chose to dichotomise our outcome variable due to the skewed distribution of our outcome data, and based on consensus between the research team and community partners. This may have reduced some of the variability in the results, but it allowed for more meaningful and intuitive public health messages to be drawn from the analysis as it negated the need for a log (or other) transformation to be undertaken. Additionally, the cut-off values used for our lower knowledge variables were somewhat arbitrary and were chosen to ensure sufficient sample size in each group to detect possible true associations and to minimise the risk of a type II error rate. Finally, the survey was of cross-sectional design, and we cannot draw any temporal relationships between observed associations. We reported our findings in accordance with the Strengthening the Reporting of Observational studies in Epidemiology statement for cross-sectional studies (online supplemental file 2).

## CONCLUSION

While the majority of MSM in Ireland were found to have high levels of knowledge about HIV/STI transmission, testing and treatment, our study highlighted that certain groups of MSM had knowledge deficits. We identified that lower knowledge was especially apparent among younger MSM, those MSM who had never tested for HIV and those MSM who were out to fewer people. These groups would likely benefit from targeted interventions to improve their knowledge of HIV and STIs. MSM who visited the national MSM-specific sexual health promotion

## Key messages

- ▶ Large numbers of MSM in Ireland had poor knowledge regarding HIV and STI transmission, testing and treatment.
- ▶ Low knowledge was more apparent among younger MSM, those who had never tested for HIV, and among MSM who were out to fewer people.
- ▶ MSM who visited a national MSM-specific sexual health promotion website displayed higher knowledge levels of HIV and STI risks.
- ▶ MSM who have poorer knowledge of HIV/STI transmission pathways and testing opportunities may benefit from targeted interventions.

website also had higher knowledge levels than those who did not.

**Handling editor** Jamie Scott Frankis

**Acknowledgements** The authors wish to thank all the men who gave their time to participate in MSM Internet Survey Ireland 2015. The authors acknowledge the support of partner organisations, including the Gay Men's Health Service, Gay Health Network, HSE Sexual Health and Crisis Pregnancy Programme, HIV Ireland, and all those who helped to promote the survey. We acknowledge in particular the help and support of Coralie Giese, Maeve O'Brien, Siobhan O'Dea, Marzena Sekular, Susan Donlon, Michelle Merrigan and Fiona Lyons.

**Contributors** CC, PB and DI designed the study. CC analysed the data under the supervision of PB and DI. CC wrote the initial manuscript. All authors were involved in revising the manuscript for intellectual content, and all authors approved the final version of the manuscript.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** None declared.

**Patient consent for publication** Not required.

**Ethics approval** Ethical approval was granted by the Royal College of Physicians in Ireland (ID: RECSAF30). All participants provided informed consent, and could withdraw from the study at any time.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** Data may be obtained from a third party and are not publicly available. Currently, the unpublished MISI data are available from the Health Protection Surveillance Centre in Dublin, Ireland. It is planned that the entire dataset will be uploaded to the Irish Social Science Data archive (<https://www.ucd.ie/issda/>) in the near future.

#### ORCID iD

Cian Carey <http://orcid.org/0000-0002-9637-9603>

#### REFERENCES

- O'Leary D. The syndemic of AIDS and STDs among MSM. *Linacre Q* 2014;81:12–37.
- HPSC. *HIV in Ireland: 2018 provisional data including latest trends Dublin*. Ireland: Health Protection Surveillance Centre, 2019.
- HPSC. *Sexually transmitted infections (STIs) in Ireland: 2018 provisional data Dublin*. Ireland: Health Protection Surveillance Centre, 2019.
- ECDC. *Who regional office for Europe. HIV/AIDS surveillance in Europe 2018 – 2017 data*. Copenhagen: European Centre for Disease Prevention and Control, 2019.
- CDC. *Hiv/Aids testing Atlanta, GA: centers for disease control and prevention, 2018*. Available: <https://www.cdc.gov/hiv/basics/testing.html>
- Paz-Bailey G, Mendoza MCB, Finlayson T, et al. Trends in condom use among MSM in the United States: the role of antiretroviral therapy and seroadaptive strategies. *AIDS* 2016;30:1985–90.
- Glick SN, Morris M, Foxman B, et al. A comparison of sexual behavior patterns among men who have sex with men and heterosexual men and women. *J Acquir Immune Defic Syndr* 2012;60:83–90.
- Shaw GM, Hunter E. HIV transmission. *Cold Spring Harb Perspect Med* 2012;2:a006965.
- Hatzenbuehler ML, O'Cleirigh C, Mayer KH, et al. Prospective associations between HIV-related stigma, transmission risk behaviors, and adverse mental health outcomes in men who have sex with men. *Ann Behav Med* 2011;42:227–34.
- Beyrer C, Baral SD, van Griensven F, et al. Global epidemiology of HIV infection in men who have sex with men. *Lancet* 2012.
- Carter A, Lachowsky N, Rich A, et al. Gay and bisexual men's awareness and knowledge of treatment as prevention: findings from the momentum health study in Vancouver, Canada. *J Int AIDS Soc* 2015;18:20039.
- Hicks G, Barragan M, Franco-Paredes C, et al. Health literacy is a predictor of HIV/AIDS knowledge. *Fam Med* 2006;38:717–23.
- Knight R, Karamouzian M, Salway T, et al. Online interventions to address HIV and other sexually transmitted and blood-borne infections among young gay, bisexual and other men who have sex with men: a systematic review. *J Int AIDS Soc* 2017;20:e25017.
- Mor Z, Turner D, Livnat Y, et al. HIV infected men who have sex with men in Israel: knowledge, attitudes and sexual behavior. *BMC Infect Dis* 2017;17:679.
- Mor Z, Grayeb E, Beany A. Arab men who have sex with men in Israel: knowledge, attitudes and sexual practices. *HIV Med* 2016;17:298–304.
- Gardner C, O'Donoghue R, Dermody A. *Findings from the National HIV knowledge and attitudes survey 2017 and people living with HIV stigma survey Dublin*. HIV in Ireland, 2017.
- O'Donnell K, Fitzgerald M, Barrett P, et al. *MISI 2015 findings from the men who have sex with men Internet survey*. Dublin: Health Protection Surveillance Centre, 2016.
- Barrett P, O'Donnell K, Fitzgerald M, et al. Drug use among men who have sex with men in Ireland: prevalence and associated factors from a national online survey. *Int J Drug Policy* 2019;64:5–12.
- O'Donnell K, Fitzgerald M, Quinlan M, et al. Inequalities in HIV testing uptake and needs among men who have sex with men living in Ireland: findings from an Internet survey. *HIV Med* 2019;20:157–63.
- Card KG, Armstrong HL, Lachowsky NJ, et al. Belief in treatment as prevention and its relationship to HIV status and behavioral risk. *J Acquir Immune Defic Syndr* 2018;77:8–16.
- DiFranceisco W, Pinkerton SD, Dyatlov RV, et al. Evidence of a brief surge in safer sex practices after HIV testing among a sample of high-risk men and women. *J Acquir Immune Defic Syndr* 2005;39:606–12.
- Mutch AJ, Lui C-W, Dean J, et al. Increasing HIV testing among hard-to-reach groups: examination of rapid, a community-based testing service in Queensland, Australia. *BMC Health Serv Res* 2017;17:310.
- Davoren MP, Shiely F, Byrne M, et al. Hazardous alcohol consumption among university students in Ireland: a cross-sectional study. *BMJ Open* 2015;5:e006045.
- Scott-Sheldon LAJ, Carey MP, Carey KB. Alcohol and risky sexual behavior among heavy drinking college students. *AIDS Behav* 2010;14:845–53.
- Mair C, Ponicki WR, Gruenewald PJ. Reducing risky sex among college students: prospects for context-specific interventions. *AIDS Behav* 2016;20 Suppl 1:109–18.
- Randolph ME, Torres H, Gore-Felton C, et al. Alcohol use and sexual risk behavior among college students: understanding gender and ethnic differences. *Am J Drug Alcohol Abuse* 2009;35:80–4.
- Aggleton P, Bhana D, Clarke DJ, et al. HIV education: reflections on the past, priorities for the future. *AIDS Educ Prev* 2018;30:254–66.
- Qiao S, Zhou G, Li X. Disclosure of same-sex behaviors to health-care providers and uptake of HIV testing for men who have sex with men: a systematic review. *Am J Mens Health* 2018;12:1197–214.
- Higgins A, Doyle L, Downes C, et al. *The LGBTIreland report: national study of the mental health and wellbeing of lesbian, gay, bisexual, transgender and intersex people in Ireland*. Dublin, 2016.
- den Daas C, Döppen M, Schmidt AJ, et al. Determinants of never having tested for HIV among MSM in the Netherlands. *BMJ Open* 2016;6:e009480.

## SUPPLEMENTARY FILE

### Supplementary Methods

#### Covariate Variables

We included the following demographic covariates: age group (18-24, 25-39, 40+ years), country of birth (born in Ireland, born outside Ireland), area of residence (Dublin, outside Dublin), education level (degree level or higher, lower than degree level), employment status (employed, unemployed, student, other). Other variables included: sexual identity (gay, bisexual, straight/other); outness (out to all/almost all, more than half, less than half, few, none); HIV testing history (tested positive, last test negative less than 12 months ago, last test negative more than 12 months ago, never tested); history of an STI in the last 12 months (no/don't know, yes) and knowledge of HIV status (sure of status, unsure of status).



### Infographics

#### Infographic 1: Hepatitis B



#### Infographic 2: Gonorrhoea



**Infographic 3: Condom use****Infographic 4: HIV/STI testing**



## Supplementary Results

### Secondary Outcome Measure

The secondary analysis included 3,017 participants who answered all six questions about HIV/STI transmission (Table 4). From the multivariable analysis, significantly lower levels of knowledge were identified among those aged 18-24 years (aOR 1.41 95%CI 1.09-1.82), those educated below degree level (aOR 1.37 95%CI 1.16-1.62), those born in Ireland (aOR 1.60 95%CI 1.26-2.02) and students (aOR 1.35 95%CI 1.07-1.71). Furthermore, lower knowledge was identified among those who were unsure of their HIV status (aOR 1.31 95%CI 1.11-1.55), those out to few of their contacts (aOR 1.29 95%CI 1.01-1.64), those who never tested for HIV (aOR 2.01 95%CI 1.64-2.345) and those who had never visited the Man2Man health promotion website (aOR 1.86 95%CI 1.53-2.27).

## Supplementary Tables

**Supplemental Table 1:** Breakdown of responses for each knowledge statement

Statement Number	I Knew this already	I was not sure about this	I did not know this already	I did not know this already	I do not understand this
1	2823	57	12	7	6
2	2794	57	20	12	22
3	2839	44	11	2	9
4	2354	325	199	5	22
5	2706	142	22	3	32
6	2723	134	38	4	6
7	1970	523	284	19	109
8	2232	457	151	3	62
9	2391	385	115	1	13
10	2745	110	37	5	8
11	2387	367	140	4	7
12	1587	549	723	19	27
13	1865	598	397	13	32

**Supplemental Table 2:** Number of men knowing each number of knowledge statements

Number of Statements known	Number of Men (%)	Cumulative %
0	3 (<1)	<1
1	6 (<1)	<1
2	8 (<1)	<1
3	17 (<1)	1
4	21 (<1)	2
5	38 (1)	3
6	64 (2)	5
7	108 (4)	9
8	156 (5)	14
9	281 (10)	24
10	353 (12)	36
11	461 (16)	52
12	506 (17)	70
13	883 (30)	100

**Supplemental Table 3:** Univariable and multivariable analysis of lower knowledge of HIV and STI transmission among MSM in Ireland, MISI 2015

Number with Lower Knowledge (%)		Univariable Analysis 95% Confidence			Multivariable Analysis 95% Confidence		
		OR	Interval	P Value	aOR	Interval	P Value
<b>Age group (years)</b>							
>40	305 (37)	1	Ref	-	1	Ref	-
25-39	472 (38)	1.03	(0.86-1.24)	0.731	1.17	(0.95-1.43)	0.137
18-24	515 (55)	2.05	(1.69-2.48)	<0.001	<b>1.41</b>	<b>(1.09-1.82)</b>	<b>0.009</b>
<b>Education</b>							
Degree or higher	575 (36)	1	Ref	-	1	Ref	-
Less than degree	685 (50)	1.80	(1.55-2.08)	<0.001	<b>1.37</b>	<b>(1.16-1.62)</b>	<b>&lt;0.001</b>
<b>Area of Residence *</b>							
Dublin	537 (39)	1	Ref	-	1	Ref	-
Outside Dublin	679 (47)	1.39	(1.20-1.62)	<0.001	1.04	(0.88-1.23)	0.633
Missing	76 (37)	0.92	(0.68-1.24)	0.582	0.81	(0.59-1.13)	0.214
<b>Employment</b>							
Employed	731 (38)	1	Ref	-	1	Ref	-
Unemployed	100 (50)	1.63	(1.22-2.18)	0.001	1.29	(0.94-1.76)	0.117
Student	374 (55)	2.07	(1.73-2.47)	<0.001	<b>1.35</b>	<b>(1.07-1.71)</b>	<b>0.01</b>
Other <sup>ψ</sup>	56 (44)	1.29	(0.90-1.86)	0.161	1.35	(0.91-2.00)	0.136
<b>Country Born</b>							
Outside Ireland	136 (32)	1	Ref	-	1	Ref	-
Ireland	1150 (45)	1.70	(1.37-2.12)	<0.001	<b>1.60</b>	<b>(1.26-2.02)</b>	<b>&lt;0.001</b>
<b>Sexual Identity</b>							
Gay	963 (41)	1	Ref	-	1	Ref	-
Bisexual	196 (49)	1.39	(1.06-1.83)	0.018	0.91	(0.70-1.18)	0.487
Straight/Other	111 (49)	1.59	(0.86-2.92)	0.139	0.87	(0.64-1.18)	0.368
<b>Outness Level</b>							
Out to all/almost all	569 (38)	1	Ref	-	1	Ref	-
More than Half	196 (45)	1.34	(1.08-1.66)	0.008	1.19	(0.94-1.49)	0.149
Less than Half	111 (42)	1.20	(0.92-1.57)	0.177	0.99	(0.74-1.33)	0.968
Few	236 (50)	1.64	(1.34-2.03)	<0.001	<b>1.29</b>	<b>(1.01-1.64)</b>	<b>0.04</b>
None	140 (54)	1.94	(1.48-2.52)	<0.001	1.34	(0.97-1.85)	0.073
<b>HIV Testing History</b>							
Negative <12 months	374 (33)	1	Ref	-	1	Ref	-
Negative >12 months	226 (37)				1.08	(0.86-1.34)	0.505
Positive	26 (17)	0.41	(0.26-0.63)	<0.001	<b>0.49</b>	<b>(0.31-0.77)</b>	<b>0.002</b>
Never Tested	655 (60)	2.88	(2.46-3.36)	<0.001	<b>2.01</b>	<b>(1.64-2.35)</b>	<b>&lt;0.001</b>
<b>STI in last 12m</b>							
No/Don't Know	1200 (44)	1	Ref	-	1	Ref	-
Yes	84 (30)	0.53	(0.41-0.69)	<0.001	0.85	(0.63-1.13)	0.261
<b>Certainty re HIV Status</b>							
Sure of HIV Status	803 (40)	1	Ref	-	1	Ref	-
Unsure of HIV Status	480 (49)	1.46	(1.25-1.70)	<0.001	<b>1.31</b>	<b>(1.11-1.55)</b>	<b>0.001</b>
<b>Visited Man2Man Website</b>							
Visited	230 (29)	1	Ref	-	1	Ref	-
Not visited	1034 (47)	2.16	(1.81-2.57)	<0.001	<b>1.86</b>	<b>(1.53-2.27)</b>	<b>&lt;0.001</b>
<b>Viewed one or more HP Infographics+</b>							
Viewed	584 (38)	1	Ref	-	1	Ref	-
Not Viewed	668 (47)	1.44	(1.25-1.67)	<0.001	1.17	(0.99-1.39)	0.063

\*Variable with &gt;5% missing data

<sup>ψ</sup> Other includes: retired; sick leave; medically retired & other

+HP, health promotion

**Supplemental Table 4:** Univariable and multivariable sensitivity analysis of lower knowledge of HIV and STI transmission, testing and treatment among HIV negative MSM in Ireland, MISI 2015

Number with Lower Knowledge (%)		Univariable Analysis 95% Confidence Interval P Value			Multivariable Analysis 95% Confidence Interval P Value		
		OR			aOR		
<b>Age group (years)</b>							<0.001
>40	226 (31)	1	Ref	-	1	Ref	-
25-39	350 (31)	0.97	(0.79 – 1.19)	0.77	<b>1.27</b>	<b>(1.01 – 1.60)</b>	<b>0.04</b>
18-24	466 (51)	2.32	(1.89 – 2.85)	<0.001	<b>2.02</b>	<b>(1.52 – 2.67)</b>	<b>&lt;0.001</b>
<b>Education</b>							<0.001
Degree or higher	422 (29)	1	Ref	-	1	Ref	-
Less than a degree	597 (48)	2.23	(1.90 – 2.61)	<0.001	<b>1.62</b>	<b>(1.35 – 1.93)</b>	<b>&lt;0.001</b>
<b>Area of Residence *</b>							0.072
Dublin	397 (32)	1	Ref	-	1	Ref	-
Outside Dublin	584 (44)	1.68	(1.43 – 1.97)	<0.001	<b>1.20</b>	<b>(1.00 – 1.44)</b>	<b>0.042</b>
Missing	61 (36)	1.20	(0.86 – 1.67)	0.29	1.02	(0.71 – 1.47)	0.90
<b>Employment</b>							0.14
Employed	570 (32)	1	Ref	-	1	Ref	-
Unemployed	81 (46)	1.77	(1.29 – 2.41)	<0.001	1.20	(0.85 – 1.69)	0.30
Student	321 (50)	2.10	(1.75 – 2.52)	<0.001	1.10	(0.86 – 1.42)	0.43
Other <sup>ψ</sup>	49 (43)	1.62	(1.10 – 2.38)	0.01	<b>1.67</b>	<b>(1.09 – 2.55)</b>	<b>0.019</b>
<b>Country of Birth</b>							<0.001
Outside Ireland	104 (27)	1	Ref	-	1	Ref	-
Ireland	933 (40)	1.78	(1.40 – 2.26)	<0.001	<b>1.62</b>	<b>(1.25 – 2.11)</b>	<b>&lt;0.001</b>
<b>Sexual Identity</b>							0.72
Gay	748 (35)	1	Ref	-	1	Ref	-
Bisexual	184 (48)	1.70	(1.37 – 2.12)	<0.001	1.01	(0.77 – 1.32)	0.96
Straight/Other	95 (45)	1.53	(1.15 – 2.04)	0.004	0.83	(0.59 – 1.16)	0.27
<b>Outness Level</b>							0.004
Out to all/almost all	425 (32)	1	Ref	-	1	Ref	-
More than Half	153 (38)	1.31	(1.04 – 1.66)	0.02	1.15	(0.89 – 1.48)	0.29
Less than Half	92 (37)	1.29	(0.97 – 1.71)	0.08	1.03	(0.76 – 1.42)	0.84
Few	206 (46)	1.88	(1.51 – 2.34)	<0.001	<b>1.43</b>	<b>(1.10 – 1.85)</b>	<b>0.007</b>
None	136 (54)	2.53	(1.92 – 3.32)	<0.001	<b>1.76</b>	<b>(1.26 – 2.47)</b>	<b>&lt;0.001</b>
<b>HIV Testing History</b>							<0.001
Negative in < 12 months	267 (25)	1	Ref	-	1	Ref	-
Negative in > 12 months	166 (28)	1.22	(0.98 – 1.54)	0.08	1.14	(0.89 – 1.46)	0.30
Never Tested	598 (56)	3.96	(3.30 – 4.76)	<0.001	<b>2.30</b>	<b>(1.86 – 2.85)</b>	<b>&lt;0.001</b>
<b>STI in last 12m</b>							0.63
No/Don't Know	978 (39)	1	Ref	-	1	Ref	-
Yes	57 (24)	0.49	(0.36 – 0.67)	<0.001	0.85	(0.60 – 1.19)	0.33
<b>Certainty of HIV Status</b>							0.001
Sure of HIV Status	636 (35)	1	Ref	-	1	Ref	-
Unsure of HIV Status	399 (43)	1.38	(1.17 – 1.62)	<0.001	<b>1.34</b>	<b>(1.12 – 1.60)</b>	<b>0.001</b>
<b>Visited Man2Man Website</b>							<0.001
Visited	173 (24)	1	Ref	-	1	Ref	-
Not visited	847 (42)	2.28	(1.88 – 2.76)	<0.001	<b>1.87</b>	<b>(1.50 – 2.33)</b>	<b>&lt;0.001</b>
<b>Viewed one or more HP Infographics+</b>							0.07
Viewed	460 (33)	1	Ref	-	1	Ref	-
Not Viewed	545 (42)	1.46	(1.25 – 1.71)	<0.001	1.16	(0.96 – 1.39)	0.12

\*Variable with &gt;5% missing data

<sup>ψ</sup> Other includes: retired; sick leave; medically retired & other

+HP, health promotion



**Supplemental Table 5:** Baseline characteristics of study sample and excluded men

<b>Age</b>		
18-24	914 (31)	51 (28)
25-39	1205 (42)	71 (38)
>40	786 (27)	63 (34)
<b>Education</b>		
Degree	1541 (53)	85 (46)
Less than a Degree	1306 (45)	91 (49)
Missing	58 (2)	9 (5)
<b>Employment</b>		
Employed	1882 (65)	110 (59)
Unemployed	190 (7)	15 (8)
Student	653 (22)	40 (22)
Other <sup>ψ</sup>	122 (4)	12 (7)
Missing	52 (2)	8 (4)
<b>Area of Residence *</b>		
Dublin	1329 (46)	67 (36)
Outside Dublin	1392 (48)	87 (47)
Missing	184 (6)	31 (17)
<b>Country of Birth</b>		
Ireland	2478 (85)	161 (87)
Outside Ireland	413 (14)	17 (9)
Missing	14 (<1)	7 (4)
<b>Sexual Identity</b>		
Gay	2261 (78)	150 (81)
Bisexual	389 (13)	17 (9)
Straight/Other	217 (8)	13 (7)
Missing	38 (1)	5 (3)
<b>Outness Level</b>		
Out to all/Almost all	1439 (50)	92 (50)
More than half	421 (14)	28 (15)
Less than half	259 (9)	15 (8)
Few	463 (16)	19 (10)
None	253 (9)	14 (8)
Missing	70 (2)	17 (9)
<b>HIV Status</b>		
HIV positive	142 (5)	10 (5)
HIV negative <12m	1088 (38)	75 (41)
HIV negative >12m	592 (20)	34 (18)
Never Tested	1062 (36)	61 (33)
Missing	21 (1)	5 (3)
<b>STI in last 12m</b>		
No/Don't Know	2611 (90)	162 (87)
Yes	273 (9)	16 (9)
Missing	21 (1)	7 (4)
<b>Certainty of HIV Status</b>		
Sure of Status	938 (32)	72 (39)
Unsure of Status	1953 (68)	111 (60)
Missing	14 (<1)	2 (1)
<b>Visited Man2Man</b>		
Visited	757 (26)	41 (22)
Not visited	2099 (72)	132 (71)
Missing	49 (2)	12 (7)
<b>Viewed HP Infographics</b>		
Viewed	1466 (50)	96 (52)
Not Viewed	1361 (47)	74 (40)
Missing	78 (3)	15 (8)

We excluded 185 men who did not answer all 13 questions relating to HIV/STI treatment, transmission and testing

\*Variable with >5% missing data

<sup>ψ</sup> Other includes: retired; sick leave; medically retired & other

STROBE Statement—Checklist of items that should be included in reports of *cross-sectional studies*

Item Number	Section	Recommendation	Presence in Study	Page Number
1	Title and abstract	(a) Indicate the study's design with a commonly used term in the title or the abstract	Y	2
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	Y	2
	Introduction			
2	Background/rationale	Explain the scientific background and rationale for the investigation being reported	Y	3
3	Objectives	State specific objectives, including any prespecified hypotheses	Y	3
	Methods			
4	Study design	Present key elements of study design early in the paper	Y	4
5	Setting	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	Y	4
6	Participants	Give the eligibility criteria, and the sources and methods of selection of participants	Y	4
7	Variables	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	Y	4 & 5
8	Data sources/measurement	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	Y	4-6
9	Bias	Describe any efforts to address potential sources of bias	Y	4
10	Study size	Explain how the study size was arrived at	Y	4
11	Quantitative variables	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	Y	6
12	Statistical methods	(a) Describe all statistical methods, including those used to control for confounding	Y	6
		(b) Describe any methods used to examine subgroups and interactions	Y	6
		(c) Explain how missing data were addressed	Y	6
		(d) If applicable, describe analytical methods taking account of sampling strategy	Y	6
		(e) Describe any sensitivity analyses		
	Results			
13	Participants	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	Y	8
		(b) Give reasons for non-participation at each stage	N	-
		(c) Consider use of a flow diagram	N	-
14	Descriptive data	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders (b) Indicate number of participants with missing data for each variable of interest	Y N	8 -
15	Outcome data	Report numbers of outcome events or summary measures	Y	8
16	Main results	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	Y	10
		(b) Report category boundaries when continuous variables were categorized	Y	10
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	N/A	-
17	Other analyses	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	Y	11
	Discussion			
18	Key results	Summarise key results with reference to study objectives	Y	13
19	Limitations	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	Y	14 & 15
20	Interpretation	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	Y	12 - 15
21	Generalisability	Discuss the generalisability (external validity) of the study results	Y	15
	Other information			
22	Funding	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	Y	15