P246 THE ROLE OF PRIOR SYPHILIS IN THE CLINICAL PRESENTATION OF THIS DISEASE AMONG ADULTS FROM STI CLINICS IN PERU

¹E Reyes Diaz*, ^{1,2}K Konda, ¹S Vargas, ¹G Calvo, ¹C Cáceres, ²J Klausner. ¹Centro de Investigación Interdisciplinaria en Sexualidad, SIDA y Sociedad (CISSS), UPCH, Lima, Perú; ²University of California, Los Angeles (UCLA), Los Angeles, USA

10.1136/sextrans-2021-sti.327

Background Individuals with prior syphilis are believed to be less symptomatic at subsequent infections. We present preliminary data using baseline information from the PICASSO cohort among people with active syphilis to compare the clinical presentations of individuals with and without prior *Treponema Pallidum* (TP) infection.

Methodology We included people with active early syphilis who fulfilled one of the following criteria: 1) current RPR titer of 1:8 plus a four-fold rise in RPR results within a year or 2) a new lesion with a positive TP-PCR DNA result from lesion swab. Syphilis naive (first infection) was defined as having a previous non-reactive TP-rapid test (TP-RT) within a year, and reinfection as having a prior reactive TP-RT or a prior RPR $\geq 1:8$ within a year. Chi2-square tests were used to examine associations.

Results Among the 91 men and transgender women with early syphilis: there were 10 primary (11%), 10 secondary (11%) and 71 latent or asymptomatic (78%) infections. Penile painless ulcers were the most common presentation of primary syphilis (80%, 8/10). All secondary syphilis cases had cutaneous rash. Thirty-six (40%) were syphilis naive, and 62 (68%) were HIV negative. Of the latter, 48/62 (77%) were on PrEP. Symptomatic syphilis was more frequent in the syphilis naive group (39% vs 11%, chi2 $p=0.002$). Among 62 HIV-negative participants, symptomatic syphilis was less frequent among PrEP users (15% vs 43%, chi2 $p=0.022$). Prior syphilis was also higher among PrEP users but no statistically significant (54% vs 27%, chi2 $p=0.063$). No significant difference was found by HIV status.

Conclusion Prior TP infection impacts clinical presentation during subsequent infections, reducing the likelihood of diagnosing primary or secondary syphilis. The finding of less frequent symptomatic cases among PrEP users may be related to higher prior syphilis among them in our study.

P250 POTENTIAL INTERACTIONS BETWEEN THE PATHWAYS TO DIAGNOSIS OF HIV/STI AND HIV SELF-TESTING: A QUALITATIVE STUDY OF GAY MEN IN SINGAPORE

¹R Tan*, ²Y Chan, ²M Ibrahim, ²L Ho, ²O Lim, ²B Choong, ³M Chio, ²M Chen, ⁴C Wong. ¹Saw Swee Hock School of Public Health, National University of Singapore, Singapore, Singapore; ²National Centre for Infectious Diseases, Singapore, Singapore; ³Department of STI Control (DSC) Clinic, National Skin Centre, Singapore, Singapore; ⁴Global Health and Population Research, FHI 360, Durham, USA

10.1136/sextrans-2021-sti.328

Background This study draws on qualitative insights on the barriers and facilitators to HIV testing, as well as perceptions of HIV self-testing (HIVST), to propose a framework to understand the benefits, but also potential knock-on implications of introducing HIVST in the context of other STI testing.

Methods We conducted semi-structured, in-depth interviews with 30 gay, bisexual and other men who have sex with men (GBMSM) aged 18 and 39 years old in Singapore. Interview

topics included barriers and facilitators to HIV and other STI testing, as well as perceptions of HIVST. Interviews were audio-recorded, transcribed, coded, and analysed using thematic analysis.

Results For HIV testing, participants cited the perceived risk of acquiring, susceptibility to, and symptoms of HIV as internal motivators, while social influence and accessibility of HIV testing services were external motivators. For STI testing, perceived symptoms and partner notification of STI were reported as an internal and external motivator, respectively. Availability of bundle tests, starting a new relationship, and instances of mandatory testing motivated both simultaneous HIV and other STI testing. The fear of a positive diagnosis and lack of confidentiality were cited as internal and external barriers to HIV testing, respectively, while low perceived severity of other STI and the cost of STI tests were cited as internal and external barriers to other STI testing, respectively. We identified pathways to HIV and other STI testing and discussed how the introduction of HIVST may reduce opportunities for other STI testing.

Conclusions The findings of this study suggest that introducing HIVST might weaken linkages to other STI testing if alternative strategies of promoting other STI testing are not simultaneously implemented. We recommend that future interventions address both the risks of HIV and other STI simultaneously, and that structural interventions promoting HIV and other STI prevention be balanced accordingly.

P254 THE LASTING IMPACT OF THE COVID 19 PANDEMIC ON SEXUAL HEALTH CARE IN A METROPOLITAN REGION

¹A Trolard*, ²L Weingarten, ¹J Cherabie, ¹H Reno. ¹Washington University, Saint Louis, USA; ²Saint Louis County Health Department, Saint Louis, USA

10.1136/sextrans-2021-sti.329

Background In the US, health policies accompanying the COVID-19 pandemic led to restricted healthcare access and decreased clinical services. Low public health funding was a barrier to accessing sexual healthcare before the pandemic and intensified during restrictions. Surveillance of the sexual healthcare landscape can inform program development to improve sexual health.

Methods We collected information from clinics offering STI testing in the St. Louis region, Missouri, USA twice during 2020: in the spring during the first pandemic lockdown and in the fall as restrictions eased. Clinics were interviewed about sexual health services and categorized into three categories: open fully with no changes; modified if hours open and/or the ability to accept walk-ins were reduced; or fully closed. We also collected chlamydia rates by zip code and categorized zip codes as low (0–174 cases per 100,000); medium (175–653); or high (654–1291) prevalence.

Results Of the 112 clinics, 47 are federally qualified health centers (serve the un- and underinsured); 16 were local public health authorities; and the remaining 49 include community, school, and private settings. In the fall, 29% of clinics were open, 55% modified, 5% closed compared with spring when 7% of clinics were open, 63% were modified, 17% were closed. There was a 450% and 200% increase in the number of clinics that were open in high and medium prevalence zip codes respectively in the fall compared to the spring. Forty-nine clinics remain modified or closed in medium and high prevalence zip codes.