

Supplemental Table: Characteristics of included and excluded participants in the analysis of chlamydia and gonorrhea testing, OHTN Cohort Study, 2008-2011

	Included Participants (N=3165)		Excluded Participants (N=1030)		P-value
	N	%	N	%	
Mean age at baseline (SD) ^a		45.5 (10.0)		46.8 (10.3)	0.0003
STI Group					
MSM	2179	68.9%	550	53.4%	<.0001
Non-MSM male	383	12.1%	251	24.4%	
Female	534	16.9%	203	19.7%	
Unknown	69	2.2%	26	2.5%	
Median year of HIV diagnosis (IQR)		1998 (1992-2004)		1997 (1991-2003)	0.05
Region of Ontario					
Toronto	2642	83.5%	0	0.0%	<.0001
Other	523	16.5%	1030	100.0%	
Ethnicity					
White	1953	61.7%	669	65.0%	<.0001
Black/African	451	14.3%	100	9.7%	
Aboriginal	142	4.5%	138	13.4%	
Other	525	16.6%	77	7.5%	
Unknown	94	3.0%	46	4.5%	
Education					
High school or less	904	28.6%	338	32.8%	<.0001
Trade school or college	921	29.1%	245	23.8%	
University	1118	35.3%	224	21.8%	
Unknown	222	7.0%	223	21.7%	
Income					
Less than \$20,000	1246	39.2%	387	37.6%	0.0006
\$20,000 to \$59,999	1086	34.3%	289	28.1%	
\$60,000 or more	538	17.0%	118	11.5%	
Unknown	295	9.3%	236	22.9%	
Median # months of prospective follow-up (IQR)		36.0 (26.4-42.0)		22.2 (12.0-37.2)	<.0001
Median CD4 cell count/mm ³ at baseline (IQR) ^{a,b}		470 (330-640)		468 (325.75-642.50)	0.726
Antiretroviral medication any time during follow-up	2937	92.8%	984	95.5%	0.002
Viral load at baseline ^a					
Undetectable (<50)	1799	56.8%	693	61.5%	<.0001
Detectable but suppressed (50-199)	250	7.9%	44	4.3%	
Unsuppressed (200+)	1079	34.1%	293	28.5%	
Unknown	37	1.2%	60	5.8%	

SD, standard deviation. IQR, interquartile range.

a) Baseline was defined as later of January 1, 2008 or enrolment date.

b) CD4 cell count was missing for 2.4% of participants for included participants and 20% for excluded participants.