

Supplementary Table: Behavioural and psychosocial characteristics and baseline demographic variables of users of daily oral PrEP (emtricitabine/tenofovir disoproxil), associated with TFV-DP concentration in linear regression analysis using generalised estimating equations. Sensitivity analysis excluding visits on which participants switched to event-driven PrEP (n=250; total number of DBS samples=440). AMPrEP study, Amsterdam, 2015-2018.

	Univariable			Multivariable		
	β^a	(95% CI)	p value	β^a	(95% CI)	p value
Demographic characteristics						
Age (modelled as cubic spline; ref.: 21 years)			0.0063			0.0014
35 years	35.4	(-248 to 319)		29.7	(-260 to 319)	
50 years	21.8	(-228 to 271)		28.5	(-226 to 283)	
55 years	131	(-120 to 382)		144	(-110 to 399)	
60 years	288	(1.06 to 576)		308	(21.5 to 595)	
65 years	469	(107 to 832)		496	(141 to 852)	
Self-declared ethnicity: non-white	-88.6	(-253 to 76.0)	0.29			
Place of residency in the Netherlands: outside Amsterdam	-31.4	(-159 to 96.3)	0.63			
Education level						
Low, middle, other	0		0.74			
High	-24.1	(-166 to 118)				
Employment ^b						
Employed	0		0.30			
Unemployed	-123	(-462 to 216)				
Other (retired, volunteer, disabled, student)	129	(-61.8 to 320)				
Net monthly income in Euro ^c						
≤1700	0		0.97			
1701-2950	44.0	(-117 to 205)				
>2950	39.5	(-132 to 211)				
Living situation						
Alone	0		0.029			
With partner	29.8	(-95.8 to 155)				
With parents/flatmates	-202	(-376 to -28.9)				
Steady relationship ^d	-17.6	(-136 to 101)	0.77			
Sexual preference: not exclusively homosexual ^e	161	(14.5 to 308)	0.031			
Sexual behaviour (past 3 months)						
Any sexually transmitted infection ^f	-32.3	(-136 to 71.2)	0.54			
Total number of sex partners (log transformed) ^g	40.1	(-9.92 to 90.1)	0.12			
Total number of condomless anal sex acts with casual partners (log transformed) ^h	25.3	(-16.1 to 66.7)	0.23			
Condomless anal sex with a casual partner (6 months prior to inclusion in AMPrEP)	191	(48.4 to 333)	0.0086	183	(50.5 to 315)	0.0068
Mental health characteristics and drug use						
Score ≥24 on Sexual compulsivity scale ^g (indication of sexual compulsivity)	-46.9	(-173 to 79.3)	0.47			
Chemsex ^{h,i}	-5.53	(-105 to 93.6)	0.91			
MHI-5 score <60 ^{g,j}	-52.3	(-186 to 81.4)	0.44			
Score ≥8 on Alcohol use disorder identification test (AUDIT) ^k	-49.2	(-171 to 73.0)	0.43			
Score ≥8 on Drug use disorder identification test (DUDIT) ^l	-42.7	(-152 to 66.9)	0.58			
Neutral to high concern about acquiring HIV ^m	104	(-70.3 to 278)	0.24			
Very important to prevent HIV ^m	83.2	(-51.9 to 218)	0.23			
Access to mobile application						
Extended app	126	(2.01 to 251)	0.046	136	(18.5 to 253)	0.023
AMPrEP study visit						
24 months vs. 12 months	-67.7	(-131 to -4.77)	0.035	-76.1	(-139 to -13.4)	0.018

AMPrEP: Amsterdam PrEP demonstration project; DBS: dried blood spots; MHI-5: Mental Health Inventory-5; PrEP: pre-exposure prophylaxis; TFV-DP: tenofovir diphosphate.

^a TFV-DP concentration in DBS in fmol/punch; ^b 6 missing; ^c 27 missing; ^d 5 missing; ^e 2 missing; ^f 4 missing; ^g 14 missing; ^h 16 missing; ⁱ Use of γ -hydroxybutyrate, γ -Butyrolactone, methamphetamine or mephedrone prior to or during sex in the 3 months prior to inclusion into AMPrEP; ^j Indication of an anxiety or depressive mood disorder; ^k Indication of an alcohol use disorder, 15 missing; ^l Indication of a drug use disorder, 15 missing; ^m Scale 1-7, dichotomised, at baseline.